



# Are Differences in the Oral Microbiome Due to Ancestry or Socioeconomics?

Christopher Kenyon<sup>a</sup>

<sup>a</sup>Department of Clinical Science, Institute of Tropical Medicine, Antwerp, Belgium

**KEYWORDS** oral microbiome, ancestry

In their recent paper, Yang et al. found differences between European Americans (EAs) and African Americans (AAs) in the abundance of various oral bacterial species based on 16S rRNA sequencing of a cohort from the southeastern United States (1). They concluded that these differences are due to intrinsic ancestry-based differences in oral microbiomes. I would advocate caution in reaching this conclusion based on the small differences identified and the concern that these differences may be due to inadequate controlling for socioeconomic factors. The differences in relative abundance of the 13 most common taxa between EA and AA were small (batch 1, median 1.2% [interquartile range {IQR}, 0.5 to 4%]; batch 2, median 0.2% [IQR, 0.2 to 0.9%]). There were large and statistically significant differences in age, gender, income, highest level of education, smoking, and proportion of participants with tooth loss between the two groups. While some of these differences were controlled for in the analyses, it is quite possible that they were inadequately controlled for. For example, only 13.1% of AAs versus 25.1% of EAs retained all of their teeth. A greater proportion of AAs had lost 1 to 10 and more than 10 teeth. This is a crude measure of periodontal health. Controlling with a more detailed measure of periodontal health may have explained most or all the differences found. Dental caries have been found to be more common in AAs than EAs and to be strongly associated with lower income and poorer education (2–4). The finding by Yang et al. that bacterial taxa that are well-known causes of periodontal disease, such as *Porphyromonas gingivalis*, *Prevotella intermedia*, *Treponema denticola*, and *Filifactor alocis*, were more prevalent in AAs than EAs is commensurate with this socioeconomic explanation.

## REFERENCES

1. Yang Y, Zheng W, Cai Q, Shrubsole MJ, Pei Z, Brucker R, Steinwandel M, Bordenstein SR, Li Z, Blot WJ, Shu X-O, Long J. 2019. Racial differences in the oral microbiome: data from low-income populations of African ancestry and European ancestry. *mSystems* 4:e00639-19. <https://doi.org/10.1128/mSystems.00639-19>.
2. Eke PI, Dye BA, Wei L, Slade GD, Thornton-Evans GO, Borgnakke WS, Taylor GW, Page RC, Beck JD, Genco RJ. 2015. Update on prevalence of periodontitis in adults in the United States: NHANES 2009 to 2012. *J Periodontol* 86:611–622. <https://doi.org/10.1902/jop.2015.140520>.
3. Dye B, Thornton-Evans G, Li X, Iafolla T. 2015. Dental caries and tooth loss in adults in the United States, 2011–2012. *NCHS Data Brief* (197):197.
4. Dye BA, Weatherspoon DJ, Mitnik GL. 2019. Tooth loss among older adults according to poverty status in the United States from 1999 through 2004 and 2009 through 2014. *J Am Dent Assoc* 150:9–23.e3. <https://doi.org/10.1016/j.adaj.2018.09.010>.

**Citation** Kenyon C. 2020. Are differences in the oral microbiome due to ancestry or socioeconomic factors? *mSystems* 5:e00836-19. <https://doi.org/10.1128/mSystems.00836-19>.

**Editor** Jian Xu, Qingdao Institute of Bioenergy and Bioprocess Technology, Chinese Academy of Sciences

**Copyright** © 2020 Kenyon. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to [ckenyon@itg.be](mailto:ckenyon@itg.be).

For the author reply, see <https://doi.org/10.1128/mSystems.00891-19>.

 Differences in oral microbiomes between African-ancestry and European-ancestry populations may be explained by socioeconomic factors and not ancestry

**Published** 10 March 2020