

Clonal propagation and the fast generation of karyotype diversity: an *in vitro* *Leishmania* model

J.-C. DUJARDIN^{1*}, S. DE DONCKER¹, D. JACQUET¹, A.-L. BAÑULS², M. BALAVOINE², D. VAN BOCKSTAELE³, M. TIBAYRENC², J. AREVALO⁴ and D. LE RAY¹

¹ Unit of Molecular Parasitology, Instituut voor Tropische Geneeskunde, 155 Nationalestraat, B-2000 Antwerpen, Belgium

² Génétique et Evolution des Maladies Infectieuses, UMR CNRS/IRD 2724, Montpellier, France

³ Department of Medicine, University Hospital of Antwerp, Wilrijkstraat 10, B-2650 Edegem, Belgium

⁴ Departamento de Bioquímica, Biología Molecular y Farmacología, Facultad de Ciencias y Filosofía and Instituto de Medicina Tropical 'Alexander von Humboldt', Universidad Peruana Cayetano Heredia, A.P. 5045, Lima 100, Peru

(Received 2 June 2006; revised 26 June 2006; accepted 26 June 2006; first published online 18 September 2006)

SUMMARY

In the present work we studied the karyotype stability during long-term *in vitro* maintenance in 3 cloned strains of *Leishmania* (*Viannia*) *peruviana*, *Leishmania* (*Viannia*) *braziliensis* and a hybrid between both species. Only the *L. (V.) peruviana* strain showed an unstable karyotype, even after subcloning. Four chromosomes were studied in detail, each of them characterized by homologous chromosomes of different size (heteromorphy). Variations in chromosome patterns during *in vitro* maintenance were rapid and discrete, involving loss of heteromorphy or appearance of additional chromosome size variants. The resulting pattern was not the same according to experimental conditions (subinoculation rate or incubation temperature), and interestingly, this was associated with differences in growth behaviour of the respective parasites. No change in total ploidy of the cells was observed by flow cytometry. We discuss several mechanisms that might account for this variation of chromosome patterns, but we favour the occurrence of aneuploidy, caused by aberrant chromosome segregation during mitosis. Our results provide insight into the generation of karyotype diversity in natural conditions and highlight the relativity of the clone concept in parasitology.

Key words: *Leishmania* (*Viannia*) *peruviana*, karyotype, hybrid, selection, ploidy, pseudo-sexuality.

INTRODUCTION

Leishmania (Kinetoplastida, Trypanosomatids) are parasitic protozoa causing a spectrum of clinical forms in humans and animals. These organisms may be considered as very successful parasites, when considering the number of different ecological niches they have colonized. *Leishmania* life-cycle occurs in 88 countries, in biotopes ranging from primary forest to xerophytic biotopes, from sylvatic to domestic environment, from lowlands to highlands. More than 50 species of sandflies are reported vectors and many mammals may act as reservoir, including edentates, monkeys, rodents, canids and humans (Shaw and Lainson, 1987). The genetic mechanisms underlying such a successful adaptation and spreading deserve attention.

Sexual recombination, a major actor of innovation and adaptive evolution, can occur in *Leishmania* as demonstrated by the observation of hybrids in wild populations (Dujardin *et al.* 1995*a*; Bañuls *et al.*

1997). However, it is likely too rare to have implications on the parasite population structure, which is essentially clonal (Tibayrenc and Ayala, 1999). Besides point mutations, *Leishmania* have developed a series of asexual mechanisms that might contribute efficiently to parasite fitness. Many genes involved in host–parasite relationships are tandemly repeated and prone to mitotic recombination, which may occur between and within tandem repeats, amplifying genes and generating diversity of genotypes (Victoir and Dujardin, 2002). Under stress conditions, the parasites can also amplify sets of genes in the form of circular amplicons or small chromosomes (Segovia and Ortiz, 1997). Last but not least, it was shown during gene knock-out experiments that *Leishmania* – normally diploid parasites (Beverley, 2003) – can undergo polyploidy and aneuploidy (Cruz *et al.* 1993). It was hypothesized that this might constitute another mechanism leading to genetic polymorphism, during reversion to the normal ploidy. For instance, a heterozygote AB changed to AABB could generate AA, AB or BB offspring during reversion, having an effect similar to recombination (Cruz *et al.* 1993). However, to our knowledge, this hypothesis was never explored experimentally in *Leishmania*.

* Corresponding author: Unit of Molecular Parasitology, Instituut voor Tropische Geneeskunde, 155 Nationalestraat, B-2000 Antwerpen, Belgium. Tel: +32 3 2476358. Fax: +32 3 2476359. E-mail: jcdujard@itg.be

The objective of the present work was to study karyotype stability during long-term *in vitro* maintenance in cloned strains of *Leishmania* (*L. (V.) peruviana*, *L. (V.) braziliensis* and a hybrid between both species) characterized by homologous chromosomes of different size (heteromorphy) or similar size (monomorphy). Heteromorphic clones were chosen because they might allow easier visualization of problems affecting chromosome assortment during segregation. One of the heteromorphic strains showed an extremely unstable karyotype even after subcloning. Our attention was focused on the latter strain, and the influence of environmental conditions as well as total and partial ploidy were analysed.

MATERIALS AND METHODS

Parasites

Three *Leishmania* cloned stocks were used in the present study: *L. (V.) braziliensis* MHOM/BR/75/M2904, *L. (V.) braziliensis/L. (V.) peruviana* hybrid MHOM/PE/91/LC1419 (Dujardin *et al.* 1995a), and *L. (V.) peruviana* MHOM/PE/84/LH78 (Dujardin *et al.* 1993b). Parasites were cultivated as promastigotes in GLSH (M2904, Le Ray, 1975) or blood agar medium (LC1419 and LH78, Tobie *et al.* 1950). Characterization of the parasites was performed by multilocus enzyme electrophoresis (MLEE), random amplified polymorphic DNA (RAPD) analysis and molecular karyotyping. For the three stocks, schemas of long-term *in vitro* maintenance were followed (see Fig. 1 for LH78) and, regularly, parasites were harvested for DNA extraction and cryostabilized in liquid nitrogen. In the case of LH78, additional subcloning was performed by the micro-drop method (Van Meirvenne *et al.* 1975), starting from cryostabilized samples. For one of these subclones (1b), 3 lines were derived and maintained under different conditions: (1bA) 26 °C, sub-inoculated 1x/week (1bB) 26 °C, sub-inoculated 3x/week, and (1bC) submitted to a succession of heat (37 °C, 5 h) and cold (4 °C, 24 h) shocks and, multiplication at 26 °C (Fig. 1). Procedures for cultivation, harvesting and preparation for orthogonal field alternating gel electrophoresis (OFAGE) have been described elsewhere (Dujardin *et al.* 1993a). For comparing growth curves of different parasite lines, the same inoculum was used, and parasites were fixed in 1% Formol and counted in a Bürker chamber.

Molecular karyotyping, hybridization and densitometry

OFAGE electrophoresis using equipment described by Dujardin *et al.* (1987) were performed on 1.5% agarose gels in 0.4 x TBE running buffer at 7–12 °C for 23 h. Resolution of the whole karyotype was




Fig. 1. Flow chart showing the history of *in vitro* maintenance of *Leishmania (Viannia) peruviana* LH78 and its different lines. R, number of sub-inoculations since the initial cloning: this counting continues even after subcloning but in that case, a second counting since subcloning is reported (numbers in parentheses).

achieved by 3 distinct runs with 45- 65- and 125-s pulses respectively. The karyotype of reference strain *L. (V.) braziliensis* M2903 was used to estimate the sizes of chromosomal bands. OFAGE-resolved chromosomes (numbered after Britto *et al.* 1998) were transferred to nylon membranes (Hybond-N, Amersham), processed and hybridized according to the manufacturer's instructions. The 4 chromosome-specific probes here used contain genomic fragments isolated from *L. (V.) braziliensis* M2904 (Dujardin *et al.* 1993b). Probes pLb-149G and pLb-168S2 are derived from pLb-149 and pLb-168 respectively, and correspond to single- or low-copy number sequences (unpublished data); pLb-149G is linked with the mini-exon genes on chromosome 2 (Kebede *et al.* 1999), while pLb-168S2 is a sequence presenting the highest identity with a portion of LM7_11b.1.Contig2 (chromosome 7 of *L. (L.) major*). Probe pLb-134Sp contains a highly conserved part of gp63 gene (Dujardin *et al.* 1994), present in multiple copies on chromosome 10. Probe pLb-22 contains a gene coding for a putative serine/threonine protein kinase (Dujardin *et al.* 1993b, and unpublished data), and is present as single copy together with rDNA genes on chromosome 27 (Inga *et al.* 1998). Probes were labelled with ³²P-dCTP, by random prime labelling. Hybridizations were performed according to the manufacturer's instructions

at 65 °C. The last wash after hybridization was performed at $1 \times$ SSC. Densitometry scanning of autoradiograms was performed in duplicate using a SHARP JX-330 scanner and the Pharmacia software Imagemaster Elite 1D programme.

Flow cytometry

Promastigotes were harvested at day 3 post-inoculation (logarithmic phase) and, prior to fixation, they were washed 3 times in ice-cold Hanks buffered salt solution HBSS (without Ca^{2+} or Mg^{2+} , addition of 1 mM EDTA). After the last centrifugation, parasites were resuspended in 30 μl of HBSS, and ice-cold 70% ethanol/5% glycerol (v/v) was added stepwise under continuous vortexing. Fixed parasites (in concentrations ranging between 10^5 and 10^6 parasites/ml) were stored at -20°C until DNA measurement. Parasites were stained with the DNA-specific YOYO-1 iodide (Molecular Probes Inc.), already applied to trypanosomes by Van den Abbeele *et al.* (1999). Fixed parasites were centrifuged and incubated in 200 μl of Tris-EDTA buffer (10 mM Tris-HCl, 1 mM EDTA, pH 7.2)/0.1% Nonidet P-40 containing 30 $\mu\text{g}/\text{ml}$ trypsin during 10 min at room temperature. Afterwards, 150 μl of TE buffer with 500 $\mu\text{g}/\text{ml}$ trypsin-inhibitor and 50 $\mu\text{g}/\text{ml}$ ribonuclease A were added. After 10 min at room temperature, 150 μl of TE buffer were added containing the YOYO-1 iodide. For the latter, a final concentration of 200 nM was obtained in a total sample volume of 500 μl . Parasites were incubated at 37 °C during 20 min. Afterwards, stained samples were stored at 4 °C and protected from the light until measurement. Fluorescence intensity from $2\text{--}20 \times 10^3$ cells was measured, digitized and stored after excitation at 488 nm using a Becton-Dickinson FACScan flow cytometer. Fluorescence signals from debris and aggregated cells were gated out using pulse shape analysis. Preliminary experiments showed that the reproducibility of the fluorescence measure was 5%.

Multilocus enzyme electrophoresis (MLEE)

MLEE experiments were performed on cellulose acetate plates (Helena Laboratories) according to the method described by Ben Abderrazak *et al.* (1993) with slight modifications (Bañuls *et al.* 2000). A total of 15 enzyme systems were performed, namely: aconitase (ACON: EC 4.2.1.3), glucose 6 phosphate dehydrogenase (G6PD: EC 1.1.1.49), glucose phosphate isomerase (GPI: EC 5.3.1.9), glutamate oxaloacetate transaminase (GOT: EC 2.6.1.1), glutamate pyruvate transaminase (ALAT: EC 2.6.1.2), isocitrate dehydrogenase (IDH: EC 1.1.1.42), malate dehydrogenase Nad^+ (MDH: EC 1.1.1.37), malate dehydrogenase NADP^+ or malic enzyme (ME: EC 1.1.1.40), mannose phosphate isomerase (MPI: EC

5.3.1.8), nucleoside hydrolases, substrate inosine and substrate deoxyinosine, I and D respectively (NHI and NHD: EC 2.4.2.*), 6 phosphogluconate dehydrogenase (6PGD: EC 1.1.1.44), and phosphoglucomutase (PGM: EC 2.7.5.1). The 13 enzyme systems used made it possible to study 14 different putative loci. Indeed 2 different loci could be distinguished with the NHI system, *Nhi* 1 and *Nhi* 2.

Random amplified polymorphic DNA (RAPD)

DNA was extracted according to the protocol communicated by Sambrook *et al.* (1989). Twelve primers of the Operon Technology kits (Alabama, California) were used: A10: GTGATCGCAG; B1: GTTTCGCTCC; F4: GGTGATCAGG, N18: GGTGAGGTCA; N20: GGTGCTCCGT; R7: ACTGGCCTGA; R13: GGACGACAAG; U2: CTGAGGTCTC; U10: ACCTCGGCAC; U12: TCACCAGCCA; U16: CTGCGCTGGA, U17: ACCTGGGGAG. For each sample and each primer the pattern reproducibility was tested twice.

RESULTS

Follow-up of M2904 and LC1419 clones

The clone of *L. (V.) braziliensis* M2904 was shown to be monomorphic for the 4 chromosomes considered (2, 430 kb; 10, 700 kb; 7, 640 kb and 27, 1300 kb), while the clone of the LC1419 hybrid was heteromorphic for each of the 4 chromosomes (2, 420/470 kb; 10, 640/700 kb; 7, 640/700 kb and 27, 1300/1150 kb). Clones of M2904 and LC1419 were maintained *in vitro* for a period of 87 (633 sub-inoculations, average frequency = $1.8 \times$ /week) and 22 (127 sub-inoculations, average frequency = $1.4 \times$ /week) months respectively. All over this period of time, no significant differences were observed in the whole molecular karyotype of the different samples (as seen after ethidium bromide staining, not shown), neither in the size of the 4 chromosomes under study: M2904 and LC1419 consistently presented monomorphic and heteromorphic chromosomes respectively. The only noticeable variation appeared in samples 5 and 6 of M2904, in which a 30-kb smaller size-variant of chromosome 10 appeared besides the 700-kb major band, but disappeared in later samples. Genotypic stability was also verified for M2904 clone, through the MLEE and RAPD analysis of 3 samples at different number of sub-inoculations (83, 292 and 323): no difference was observed (not shown).

Follow-up of LH78 clone and subclones

The clone of *L. (V.) peruviana* LH78 was also heteromorphic for the 4 chromosomes under study (see below). It was first maintained *in vitro* for a

Table 1. Variation of patterns of chromosomes 2, 10, 7 and 27 during *in vitro* maintenance of *Leishmania (Viannia) peruviana* LH78 (underlined and bold, chromosomal variants with higher hybridization intensity)

(See Fig. 1 for details on sample designation.)

| | | | | | | | | | | | | | |
|-----------------|----------------|------------|------------|------------|------------|--------------|------------|------------|------------|--------------|--------------|--------------|--|
| Sample | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | | | | |
| Sub-inoculation | 16 | 34 | 54 | 72 | 92 | 102 | | | | | | | |
| Chromosome | | | | | | | | | | | | | |
| 27 | <u>1/3/4</u> | <u>1/3</u> | <u>1/2</u> | <u>1/2</u> | <u>1/2</u> | | | | | | | | |
| 7 | 1/2 | <u>1/2</u> | <u>1/2</u> | <u>1/2</u> | <u>1/2</u> | | | | | | | | |
| 10 | 1/2 | 1/2 | 1/2 | 1/2 | <u>1/2</u> | 1 | 1 | 1 | 1 | | | | |
| 2 | <u>1/2/3/4</u> | 1/2/3 | 1/2 | 1/2 | <u>1/2</u> | 1/2 | 1/2 | 1/2 | 1/2 | | | | |
| Sample | 1a | 1b | 1c | 5d | 5e | | | | | | | | |
| Sub-inoculation | 28(2) | 28(2) | 28(2) | | | | | | | | | | |
| Chromosome | | | | | | | | | | | | | |
| 27 | 1 | 1/3 | <u>1/3</u> | <u>1/2</u> | 1/2 | | | | | | | | |
| 7 | 1/2 | 1/2 | <u>1/2</u> | | | | | | | | | | |
| 10 | 1/2 | 1/2 | 1/2 | 1/2 | 1 | | | | | | | | |
| 2 | 1/2 | 2/3 | 2/3 | 1/2 | 1/2 | | | | | | | | |
| Sample | 1bA1 | 1bA2 | 1bA3 | 1bA4 | 1bB1 | 1bB2 | 1bB3 | 1bB4 | 1bC1 | 1bC2 | 1bC3 | 1bC4 | |
| Sub-inoculation | 47(21) | 56(30) | 65(39) | 76(50) | 71(45) | 96(70) | 122(96) | 154(128) | 65(39) | 82(56) | 100(74) | 122(96) | |
| Chromosome | | | | | | | | | | | | | |
| 27 | <u>1/3</u> | <u>1/3</u> | <u>1/3</u> | <u>1/3</u> | <u>1/3</u> | <u>1/3</u> | <u>1/3</u> | <u>1/3</u> | <u>1/3</u> | <u>1/2/3</u> | <u>1/2/3</u> | <u>1/2/3</u> | |
| 7 | <u>1/2</u> | <u>1/2</u> | <u>1/2</u> | <u>1/2</u> | <u>1/2</u> | <u>1/2</u> | <u>1/2</u> | <u>1/2</u> | <u>1/2</u> | 1/2 | 1/2 | 1/2 | |
| 10 | 1 | 1 | 1 | 1 | 1/2 | 1/2 | 1/2 | 1/2 | 1/2 | 1/2 | 1/2 | 1/2 | |
| 2 | 2/3 | 2/3 | 2/3 | <u>2/3</u> | 2/3 | <u>2/3/4</u> | 2 | 2 | 2/3 | 2/3 | 2/3 | 2/3 | |

period of 16 months (127 sub-inoculations, average frequency = 2 ×/week). During this period, patterns significantly varied for 3 chromosomes, however, variation was discrete. Chromosome 27 presented 3 variants (1100, 1420 and 1520 kb, respectively variants 1, 3 and 4) in sample 1, variant 4 disappeared in sample 2, and from sample 3, variant 3 was replaced by a new 1300-kb variant (numbered 2). With respect to chromosome 10, 2 size-variants (640 and 700 kb, respectively variants 1 and 2) were initially observed; in sample 5, the hybridization intensity of variant 1 was increased, and in sample 6 and later, variant 2 was never observed. Chromosome 2 was characterized by 4 variants (400, 435, 470 and 500 kb, respectively numbered 1–4) in sample 1, variant 4 disappeared in sample 2, and from sample 3, only variants 1 and 2 were observed. In contrast, the double banding of chromosome 7 (640 and 700 kb) remained constant until the end of monitoring.

Samples of LH78 presenting more than 2 size-variants for a given chromosome (e.g. samples 1 and 2), or – in the case of doublets – strong differences in hybridization intensity of single copy probes to one variant (e.g. sample 5) deserved particular attention. Indeed, such phenomena could be due to population heterogeneity (under a diploid hypothesis) or aneuploidy. In order to distinguish between both hypotheses, subcloning was performed starting from samples 1 and 5 (see Fig. 1). This clearly indicated that both samples were heterogeneous. On the one hand, the pattern 1/3/4 of chromosome 27 in sample 1 generated patterns 1 and 1/3 in subclones 1a and

1b-c respectively. On the other hand, the pattern 1/2 of chromosome 10 (with stronger hybridization intensity for variant 1) in sample 5 gave patterns 1 and 1/2 in subclones 5d and 5e respectively.

In all subclones presenting heteromorphic chromosomes, hybridization intensity of each variant was similar, except for subclone 1c, in which hybridization intensity of variant 3 of chromosome 27 was twice as high (as measured by densitometry) than that of variant 1. The observation of this pattern only 7 passages after subcloning might be an indication of aneuploidy for chromosome 27, but could also reveal the existence of heterogeneous populations (3/0 – haploid- + 1/3) produced after subcloning. No differences were encountered in the MLEE and RAPD patterns of the 5 samples of LH78 here analysed (1, 2 and 5 from the initial clone and subclones 1b and 1c).

In order to determine whether variation of chromosome patterns (i) was a reproducible phenomenon, and (ii) could be modulated according to maintenance conditions, 3 lines were derived from subclone 1b and followed over 7 months. Significant differences were encountered according to the cultivation protocol (Table 1). Line 1bA was sub-inoculated 1 ×/week: at passage 47, chromosome 10 was already monomorphic for the small variant; at passage 76, the small variant 2 of chromosome 2 appeared to be dominant (on the basis of hybridization intensity). Line 1bB was sub-inoculated 3 ×/week: at passage 96, 3 variants of chromosome 2 were observed (addition of variant 4), and at passage 96, a monomorphic stage was reached with the presence of

variant 2 only. Finally, line 1bC was submitted to a series of heat and cold shocks (see Materials and Methods section): at passage 82, an additional variant of chromosome 27 appeared, and the tri-variant pattern remained up to the end of the experiment.

Flow cytometry

In order to determine whether variations in chromosome patterns of LH78 could have been associated with changes in total ploidy, the DNA content of subclone 1b and all stocks of the 3 derived lines A-C was measured by flow cytometry. A classical picture was observed with a major peak corresponding to parasites with 2N DNA content and a minor 4N peak, corresponding to replicating parasites. However, no significant differences were observed between the different samples (not shown).

Growth parameters

In order to have a first insight into biological changes that might accompany the differences in chromosome patterns here observed, we compared the growth curves of lines A-C of subclone 1b at the end-point of maintenance. Differences in doubling time were observed: 17.8 h for the line sub-inoculated once a week (1bA4) versus 13.3 h for the one sub-inoculated 3 times a week (1bB4). Interestingly, the line submitted to heat-cold shocks (1bC4) presented the lowest doubling time: 11.3 h.

DISCUSSION

The karyotype of *Leishmania* is considered to be generally very stable *in vitro*, as reported earlier (Giannini *et al.* 1990), and as demonstrated here for 2 out of 3 strains under observation. However, this contrasts with a very high chromosome size polymorphism in natural populations of the parasite (Dujardin *et al.* 2002). We described here a *L. (V.) peruviana* strain showing *in vitro* a similar level of karyotype instability as in the wild. The clone (LH78) was characterized by the presence of 4 heteromorphic chromosomes and we observed the rapid appearance of a mixture of parasites with rearranged chromosome patterns during *in vitro* maintenance. This phenomenon appeared again after subcloning, and thus could not be attributed to an initial cloning error. Four mechanisms might account for this variation of chromosome patterns. (1) Amplification/deletion of tandemly repeated genes is the most frequent mechanism responsible for chromosome size-variation (Victoir and Dujardin, 2002). This was shown among natural populations of *Leishmania* for 3 of the chromosomes here studied, involving genes coding for mini-exon (chromosome 2, Kebede *et al.* 1999), gp63 (chromosome 10,

Dujardin *et al.* 1994; Victoir *et al.* 1995) and rDNA (chromosome 27, Inga *et al.* 1998). However, in natural populations, size-variation of these chromosomes had been shown to be continuous (Dujardin *et al.* 1995b), which was never observed in the present case. Thus, even if chromosome variants are characterized by different copy number of repeated genes, the discrete variation of patterns here observed is more suggestive of an effect of chromosome reassortment, and the amplification/deletion hypothesis should be rejected here. (2) Another mechanism, the subject of controversy (Bastien *et al.* 1992), is the occurrence of sexual exchange. While we have previously shown that this might happen (stock LC1419 is a putative *L. (V.) braziliensis/L. (V.) peruviana* hybrid, Dujardin *et al.* 1995a), it was also suggested to be exceptional (Bañuls *et al.* 2000; Tibayrenc and Ayala, 1999). In the present case, we believe that sexual exchange was not involved in the generation of rearranged patterns. Indeed, for each chromosome, 1 or 2 segregation genotype(s) were lacking: for instance, genotype 2/2 of chromosome 10 was never observed, and chromosome 7 was only represented by genotype 1/2. (3) One might also consider a change in the total ploidy of the parasite as described previously (Cruz *et al.* 1993), followed by a return to diploidy and the 'clonal' production of segregation and recombination-like patterns. In the present case, DNA content (as measured by flow cytometry) remained constant during *in vitro* maintenance. We cannot exclude that at certain stages, a minor proportion of the parasites went polyploid for a short time, being undetectable under our experimental conditions. (4) The last mechanism would be aneuploidy, caused by aberrant chromosome segregation during mitosis. This model would better fit with our data as it concerns individual chromosomes and is coherent with the absence of recombination-like patterns. It is supported by the observation, in freshly subcloned sample 1c, of pattern 1/3 for chromosome 27, with variant 3 hybridizing twice as high to a single copy probe as variant 1. However, in this particular case, we cannot exclude the possibility that between subcloning and chromosome analysis (7 passages), a mixture (3/0, haploid and 1/3 or 1/0, haploid and 3/3) of parasites would already have occurred.

Whatever the mechanism, these data show clearly that *Leishmania* may rearrange their karyotype at a very fast rate. Interestingly, this is not the case for all stocks; indeed, the putative hybrid LC1419 (showing the same heteromorphic patterns for chromosomes 2, 10, 7 and 27 as LH78) did not show any sign of rearrangement while kept *in vitro* for a longer period. Further work should be performed in order to determine whether this discrepancy is due to intrinsic factors affecting chromosome segregation, as observed in other organisms: mutations in hsp70 (Oka *et al.* 1998), actin (Kopecka and Gabriel, 1998) and chromosome segregation genes (Xiao *et al.* 1993)

in *Saccharomyces cerevisiae*. In a recent study on *L. (L.) donovani*, a 30-kb region was reported to be involved in the mitotic stability of a linear extra chromosome, and it was hypothesized that it could contain elements participating in replication or centromeric function (Dubessay *et al.* 2001).

Our results provide some insight into the generation of karyotype diversity in natural conditions. The region of Peru, where LH78 was isolated, is characterized by parasite populations with extremely heterogeneous karyotypes (Dujardin *et al.* 1993a). In a previous study, we demonstrated that the same population could be in linkage equilibrium for chromosome size-variants (7 and 27) and in disequilibrium for RAPD markers (Bañuls *et al.* 2000). The contrast between these data was hardly explained by sexual exchanges, but more by pseudo-sexuality. According to our present work, a single stock like LH78 could spontaneously produce aberrant chromosome assortments, without affecting loci recognized by RAPD markers. At this stage, there is no evidence that the karyotype changes directly underly the observed differences in growth rate. We previously demonstrated that size-variation of 3 of the chromosomes here studied (2, 10 and 22) was associated with dosage of major genes (mini-exon, gp63 and rDNA) in natural populations of *L. (V.) braziliensis* and *L. (V.) peruviana*, with possible functional effects (Dujardin *et al.* 2002). It is equally possible that there are inducible genes whose activity was modified by the culture conditions. Nevertheless, in the case of variable selective pressure (2 different sandfly vectors, *Lutzomyia peruensis* and *Lu. verrucarum* (Villaseca *et al.* 1993), differences in altitude and temperature), the progeny of a strain like LH78 could rapidly generate a genomically heterogeneous population.

Our work has important implications concerning the relativity of the clone concept. It is obvious from data presented here that this concept is strongly limited in time, and that theoretically, the definition of clone is strictly valid for one generation only. Furthermore, the risk of selecting parasites with different karyotypes and different biological features may be increased by variations in maintenance conditions. Even if strains like LH78 are perhaps exceptional, this risk should be taken into consideration, and its potential consequences explored, in all experiments in which a 'clone' is chosen as a homogeneity criterion. This would be particularly recommended in any study on parasite phenotype, like virulence or drug resistance. A minimal precaution would be to cryostabilize large amounts of 'cloned' material and to start any biological experiment from it.

This study was financially supported by EC (Contract TS3-CT-92-0129) and the Belgian Agency for Cooperation to Development (D. G. C. D.)

REFERENCES

- Bañuls, A. L., Guerrini, F., Le Pont, F., Barrera, C., Espinel, I., Guderian, R., Echeverria, R. and Tibayrenc, M. (1997). Evidence for hybridization by multilocus enzyme electrophoresis and random amplified polymorphic DNA between *Leishmania braziliensis* and *Leishmania panamensis/guyanensis* in Ecuador. *Journal of Eukaryotic Microbiology* **44**, 408–411.
- Bañuls, A. L., Dujardin, J. C., Guerrini, F., De Doncker, S., Jacquet, D., Arevalo, J., Noel, S., Le Ray, D. and Tibayrenc, M. (2000). Is *Leishmania (Viannia) peruviana* a distinct species? A MLEE/RAPD evolutionary genetics answer. *Journal of Eukaryotic Microbiology* **47**, 197–207.
- Bastien, P., Blaineau, C. and Pagès, M. (1992). *Leishmania*: sex, lies and karyotype. *Parasitology Today* **8**, 174–177.
- Ben Abderrazak, S., Guerrini, F., Mathieu-Daude, F., Truc, P., Neubauer, K., Lewicka, K., Barnabe, C. and Tibayrenc, M. (1993). Isoenzyme electrophoresis for parasite characterization. *Methods in Molecular Biology* **21**, 361–382.
- Beverley, S. M. (2003). Protozoomics: trypanosomatid parasite genetics comes of age. *Nature Reviews Genetics* **4**, 11–19. doi: 10.1038/nrg980
- Britto, C., Ravel, C., Bastien, P., Blaineau, C., Pagès, M., Dedet, J. P. and Wincker, P. (1998). Conserved linkage groups associated with large-scale chromosomal rearrangements between Old World and New World *Leishmania* genomes. *Gene* **222**, 107–117.
- Cruz, A. K., Titus, R. and Beverley, S. M. (1993). Plasticity in chromosome number and testing of essential genes in *Leishmania* by targeting. *Proceedings of the National Academy of Sciences, USA* **90**, 1599–1603.
- Dubessay, P., Ravel, C., Bastien, P., Lignon, M. F., Ullman, B., Pagès, M. and Blaineau, C. (2001). Effect of large targeted deletions on the mitotic stability of an extra chromosome mediating drug resistance in *Leishmania*. *Nucleic Acids Research* **29**, 3231–3240.
- Dujardin, J. C., Bañuls, A. L., Llanos-Cuentas, A., Alvarez, E., De Doncker, S., Jacquet, D., Le Ray, D., Arevalo, J. and Tibayrenc, M. (1995a). Putative *Leishmania* hybrids in the Eastern Andean valley of Huanuco, Peru. *Acta Tropica* **59**, 293–307.
- Dujardin, J. C., De Doncker, S., Victoir, K., Le Ray, D., Hamers, R. and Arevalo, J. (1994). Size polymorphism of chromosomes bearing gp63 genes in *Leishmania* of the *braziliensis* complex: indication of a rearrangement of the gp63 genes in *L. braziliensis* and *L. peruviana*. *Annals of Tropical Medicine and Parasitology* **88**, 445–448.
- Dujardin, J. C., Dujardin, J. P., Tibayrenc, M., Timperman, G., De Doncker, S., Jacquet, D., Arevalo, J., Llanos-Cuentas, A., Guerra, H., Bermudez, H., Hamers, R. and Le Ray, D. (1995b). Karyotype plasticity in neotropical *Leishmania*: an index for measuring genomic distance among *L. (V.) peruviana* and *L. (V.) braziliensis* populations. *Parasitology* **110**, 21–30.
- Dujardin, J. C., Gajendran, N., Arevalo, J., Llanos-Cuentas, A., Guerra, H., Gomez, J., Arroyo, J., De Doncker, S., Jacquet, D., Hamers, R.

- and Le Ray, D. (1993 a). Karyotype polymorphism and conserved characters in the *Leishmania (Viannia) braziliensis* complex explored with chromosome-derived probes. *Annales de la Société belge de MédecineTropicale* **73**, 101–118.
- Dujardin, J. C., Gajendran, N., Hamers, R., Mathijsen, G., Urjel, R., Recacoechea, M., Villaroel, G., Bermudez, H., Desjeux, P., De Doncker, S. and Le Ray, D. (1987). Leishmaniasis in the Lowlands of Bolivia. VII. Characterization and identification of Bolivian isolates by PFG karyotyping. In *Leishmaniasis: the First Centenary (1885–1985). New Strategies for Control* (ed. Hart, D), pp. 137–148. NATO ASI series A, Plenum Press, New York.
- Dujardin, J. C., Llanos-Cuentas, A., Caceres, A., Arana, M., Dujardin, J. P., Guerrini, F., Gomez, J., Arroyo, J., De Doncker, S., Jacquet, D., Hamers, R., Guerra, H., Le Ray, D. and Arevalo, J. (1993 b). Molecular karyotype variation in *Leishmania (Viannia) peruviana*: indication of geographical populations in Peru distributed along a north-south cline. *Annals of Tropical Medicine and Parasitology* **87**, 335–347.
- Dujardin, J. C., Victoir, K., De Doncker, S., Guerbouj, S., Arevalo, J. and Le Ray, D. (2002). Molecular epidemiology and diagnosis of *Leishmania*: what have we learnt from genomic structure, dynamics and function. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **96** (S1), 81–86.
- Giannini, S. H., Curry, S., Tesh, R. and Van der Ploeg, L. H. T. (1990). Size-conserved chromosomes and stability of molecular karyotype in cloned stocks of *Leishmania major*. *Molecular and Biochemical Parasitology* **39**, 9–22.
- Inga, R., De Doncker, S., Gomez, J., Lopez, M., Garcia, R., Le Ray, D., Arevalo, J. and Dujardin, J. C. (1998). Relation between variation in copy number of ribosomal RNA encoding genes and size of harbouring chromosomes in *Leishmania* of subgenus *Viannia*. *Molecular and Biochemical Parasitology* **92**, 219–228.
- Kebede, A., De Doncker, S., Arevalo, J., Le Ray, D. and Dujardin, J. C. (1999). Size-polymorphism of mini-exon gene-bearing chromosomes among natural populations of *Leishmania*, subgenus *Viannia*. *International Journal for Parasitology* **29**, 549–557.
- Kopecka, M. and Gabriel, M. (1998). The aberrant positioning of nuclei and the microtubular cytoskeleton in *Saccharomyces cerevisiae* due to improper actin function. *Microbiology* **144**, 1783–1797.
- Le Ray, D. (1975). Structures antigéniques de *Trypanosoma brucei* (Protozoa, Kinetoplastida). Analyse immunoelectrophorétique et étude comparative. *Annales de la Société belge de MédecineTropicale* **55**, 158–160.
- Oka, M., Nakai, M., Endo, T., Lim, C. R., Kimata, Y. and Kohno, K. (1998). Loss of Hsp70-Hsp40 chaperone activity causes abnormal nuclear distribution and aberrant microtubule formation in M-phase of *Saccharomyces cerevisiae*. *Journal of Biological Chemistry* **273**, 29727–29737.
- Sambrook, J., Fritsch, E. F. and Maniatis, T. (1989). *Molecular Cloning: a Laboratory Manual*, 2nd Edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, USA.
- Segovia, M. and Ortiz, G. (1997). LD1 amplifications in *Leishmania*. *Parasitology Today* **13**, 342–348.
- Shaw, J. J. and Lainson, R. (1987). Ecology and epidemiology: New World. In *The Leishmaniasis in Biology and Medicine* (ed. Peters, W. and Killick-Kendrick, R.), vol. I, pp. 291–365. Academic Press, London.
- Tibayrenc, M. and Ayala, F. J. (1999). Evolutionary genetics of *Trypanosoma* and *Leishmania*. *Microbes and Infection* **1**, 465–472.
- Tobie, E. J., Von brand, T. and Mehlman, B. (1950). Cultural and physiological observations on *Trypanosoma rhodesiense* and *Trypanosoma gambiense*. *Journal of Parasitology* **36**, 48–54.
- Van Den Abbeele, J., Claes, Y., van Bockstaele, D., Le Ray, D. and Coosemans, M. (1999). *Trypanosoma brucei* spp. development in the tsetse fly: characterization of the post-mesocyclic stages in the foregut and proboscis. *Parasitology* **118**, 469–478.
- Van Meirvenne, N., Janssens, P. G. and Magnus, E. (1975). Antigenic variation in syringe passaged populations of *Trypanosoma (Trypanosoon) brucei*. I. Rationalization of the experimental approach. *Annales de la Société belge de MédecineTropicale* **5**, 1–23.
- Victoir, K. and Dujardin, J. C. (2002). How to succeed in parasitic life without sex? Asking *Leishmania*. *Trends in Parasitology* **18**, 81–85.
- Victoir, K., Dujardin, J. C., De Doncker, S., Barker, D. C., Arevalo, J., Hamers, R. and Le Ray, D. (1995). Plasticity of gp63 gene organization in *Leishmania (Viannia) braziliensis* and *Leishmania (Viannia) peruviana*. *Parasitology* **111**, 265–273.
- Villaseca, P., Llanos-Cuentas, A., Perez, E. and Davies, C. R. (1993). A comparative field study of the relative importance of *Lutzomyia peruensis* and *Lu. verrucarum* as vectors of cutaneous leishmaniasis in the Peruvian Andes. *American Journal of Tropical Medicine and Hygiene* **49**, 260–269.
- Xiao, Z., McGrew, J. T., Schroeder, A. J. and Fitzgerald-Hayes, M. (1993). CSE1 and CSE2, two new genes required for accurate mitotic chromosome segregation in *Saccharomyces cerevisiae*. *Molecular and Cell Biology* **13**, 4691–4702.